

Palindromic Width of Wreath Products

Elisabeth Fink[†]

19th February 2014

Abstract

We show that the wreath product $G \wr \mathbb{Z}^n$ of any finitely generated group G with $\mathbb{Z}^n$ has finite palindromic width. We also show that $C \wr A$ has finite palindromic width if C has finite commutator width and A is a finitely generated infinite abelian group. Further we prove that if H is a non-abelian group with finite palindromic width and G any finitely generated group, then every element of the subgroup $G' \wr H$ can be expressed as a product of uniformly boundedly many palindromes. From this we obtain that $P \wr H$ has finite palindromic width if P is a perfect group and further that $G \wr F$ has finite palindromic width for any finite, non-abelian group F .

1 Introduction

Palindromic words in groups have been studied from various angles lately. They make their first appearance in [Col95], where D. Collins studied palindromic automorphisms of free groups. In [GC00] H.H. Glover and C.A. Jensen study the geometry of palindromic automorphism groups of the free group. Later in [BST05] it was shown that free groups have infinite palindromic and primitive width and F. Deloup [Del10] studied the palindromic map, which is an anti-automorphism, in braid and Artin groups.

More recently, it has been established by V. Bardakov and K. Gongopadhyay [BG14c] that free nilpotent groups and free abelian-by-nilpotent groups have finite palindromic width. A paper in preparation by the same authors shows that some extensions and quotients of these groups have finite palindromic width as well [BG14a]. Another paper [BG14b] by the same authors proves that certain soluble groups have finite palindromic width. In [BG14c] they use results about the commutator width in nilpotent groups to establish their result. Independently, it has been shown by T. Riley and A. Sale in [RS14] that free metabelian groups have finite palindromic width by using results about skew-symmetric functions on free abelian groups. Further, the same authors show that $B \wr \mathbb{Z}^n$ has finite palindromic width if B is a group with finite palindromic width.

We extend this result to the case where G is any finitely generated group, then we show that $G \wr \mathbb{Z}^n$ has finite palindromic width. A result by M. Akhavan-Malayeri [AM10] shows that the wreath product of F_d with $\mathbb{Z}^n$ has finite commutator width. We use the result from [AM10] to prove that $F_d \wr \mathbb{Z}^n$ has finite palindromic width and then deduce that this property also holds for its quotients. We also give a proof that the wreath product $C \wr A$ has finite palindromic width if C is a finitely generated group that has finite commutator width and A a finitely generated infinite abelian group.

[†]This work is supported by the ERC starting grant 257110 “RaWG”

More generally, let G be any finitely generated group and H a non-abelian group which has finite palindromic width with respect to some generating set. We establish that every element of the subgroup $G' \wr H$ of the regular wreath product $G \wr H$ is a finite product of palindromes. As a Corollary we obtain that $G \wr F$ has finite palindromic width if F is a non-abelian finite group.

More concretely, we prove the following, where $pw(G, X)$ denotes the palindromic width of the group G with respect to the generating set X .

- Theorem 1.1.**
1. Let G be a d -generated group generated by X and E be the standard generating set of $\mathbb{Z}^k$, for $k \in \mathbb{N}$. Then we have that $pw(G \wr \mathbb{Z}^k, X \cup E) \leq 5d + 9k$ if k is even and $pw(G \wr \mathbb{Z}^k, X \cup E) \leq 5d + 9k + 2$ if k is odd.
 2. Assume that A is an r -generated infinite abelian group generated by T and C a finitely generated group with finite commutator width n . Then $pw(C \wr A, T \cup Y) \leq 5|Y| + 6r + 7n$, for any finite generating set Y of C .
 3. Let G be a finitely generated group and H a non-abelian finitely generated group with finite palindromic width with respect to the generating set Z . Then every element of the subgroup $G' \wr H$ can be written as a product of at most $pw(H, Z) + 1$ palindromes.
 4. Assume that P is a finitely generated perfect group and H as above. Then for any generating set S of P we have $pw(P \wr H, Z \cup \{c\} \cup S) \leq pw(H, Z) + 1$, where $c \in H$, possibly $c = 1$.
 5. If K is a non-abelian finite group and G a d -generated group, then $G \wr H$ has finite palindromic width with respect to the natural generating set.

Acknowledgement. I would like to thank Tim Riley for many helpful comments and for pointing out a crucial mistake in an earlier version of this paper.

2 Wreath Products and Words

The regular wreath product $B \wr A$ of two groups B and A is given by $F \rtimes A$, where $F = \prod_A B$ and the action of A on itself is given by left multiplication. We can write every $g \in B \wr A$ as

$$g = a \cdot \prod_{i=1}^k a_i^{-1} (f_i, 1) a_i \quad (1)$$

with $a, a_i \in A$ and $f_i \in B$ and $k \in \mathbb{Z}$. If X is a generating set for A and Y one for B , then the set $\{(y_i, 1), (1, x_i) \mid y_i \in Y, x_i \in X\}$ is a natural generating set for $B \wr A$.

A *group word* w is an element of $F_n = \langle x_1, \dots, x_n \rangle$, the free group on n generators. Examples of such words are commutators $w(x_1, x_2) = x_1^{-1} x_2^{-1} x_1 x_2$, Burnside words $w(x_1) = x_1^p$ for some $p \in \mathbb{Z}$ and many others. A word w has the general form $w(x_1, \dots, x_n) = \prod_{j=1}^k x_{i_j}^{p_j}$ for $k \in \mathbb{N}$ and $p_j \in \mathbb{Z}$. We say a word w is a *palindrome* if $x_{i_j} = x_{i_{k-j+1}}$ and $p_j = p_{k-j+1}$ for all $j = 1, \dots, k$. Examples of palindromic words are $x_1 x_2 x_1$, all power words or words of the form $x_1^2 x_2 x_1^2$. For any word w in F_n we denote by $\overline{w}$ the word

$$\overline{w} = \prod_{j=1}^k x_{i_{k-j+1}}^{p_{k-j+1}},$$

which we call the *reverse word* of w . We will use this notation throughout this paper.

Every word $w \in F_n$ defines a verbal mapping for any group G from $G^{(k)} = \prod_{i=1}^k G$ to G in a natural way. More details on this can be found in [Seg09]. We call the image

$$w(G, \dots, G) = \langle \{w(g_1, \dots, g_n) \mid g_i \in G\} \rangle$$

the *verbal subgroup* of w in G . It is a well-studied topic to decide when this group is actually generated by products of at most $C \in \mathbb{N}$ words. If there exists such a constant C , then we say that the word w has *finite width* in G . The most studied word is the commutator word, leading to the question when every element of G' can be expressed as a product of at most C commutators. Examples of groups having finite commutator width are all finite simple groups [LOST10] with commutator width 1 or most finite quasi-simple groups [LOST11]. Infinite examples are soluble groups satisfying the maximal condition on normal subgroups [AM06]. It has been shown in [Seg09, Theorem 2.1.3] that any virtually nilpotent group of finite rank has the property that any word has finite width. We note that palindromes are not group words under this definition.

However, in analogy with the terminology above, we say that an element g of a group is a palindrome, if it can be represented by a palindromic word in the generators of G . Further, a group G has *finite palindromic width* $pw(G, X)$ with respect to a generating set X , if every element of G can be expressed as a product of at most $pw(G, X)$ many palindromes in the alphabet X . If we just say a group G has finite palindromic width without reference to a generating set, then we mean that there exists a generating set with respect to which G has finite palindromic width. It is yet unclear how the choice of the generating set influences the palindromic width of G . We will however use below that if G has finite palindromic width with respect to a set X , then it has at most the same finite palindromic width with respect to any generating set containing X .

3 Wreath Product with a Free Abelian Group

In this Section we use the results of M. Akhavan-Malayeri from [AM10] and V. Bardakov and K. Gongopadhyay [BG14c] to deduce that the wreath product of a finitely generated group with a finitely generated free abelian group has finite palindromic width. We also give a self-contained proof that $C \wr A$ has finite palindromic width if C is a group with finite commutator width and A an infinite finitely generated abelian group.

We will implicitly use the following Lemma in many places. The proof of it is obvious and we therefore omit it.

Lemma 3.1. (a) *If A is an abelian group, then $a = \bar{a}$ for all $a \in A$.*

(b) *For any group G and every element $g \in G$ we have that $(\bar{g})^{-1} = \overline{g^{-1}}$.*

The following Theorem gives a uniform bound on the commutator width of a wreath product. A first, less general, version of this can also be found in M. Akhavan-Malayeri's thesis [AM97].

Theorem 3.2 ([AM10]). *Let A be a non-abelian free group and $W = A \wr B$, where B is a free abelian group of rank n . Then every element of W' is a product of at most $n + 2$ commutators, and furthermore $2 \leq cw(W) \leq n + 2$.*

We use the shape of the commutators that emerges in the proof of Theorem 3.2 to prove that the same wreath products also have finite palindromic width. The following result from [BG14c] gives an upper bound for the palindromic width of a metabelian group.

Theorem 3.3. [BG14c] *Let G be a non-abelian free abelian-by-nilpotent group of rank n . Then $pw(G, X) \leq 5n$ for a minimal generating set X .*

Combining these two Theorems allows us to obtain a bound for the palindromic width of such wreath products.

Theorem 3.4. *The wreath product $F_d \wr \mathbb{Z}^n$ of the free d -generated group F_d with $\mathbb{Z}^n$ has finite palindromic width at most $5d + 9n$ if n is even and at most $5d + 9n + 2$ if n is odd with respect to the natural generating set coming from F_d and $\mathbb{Z}^n$.*

Proof. Set $B = \prod_{\mathbb{Z}^n} F_d$ to be the base group of the wreath product. We can split the wreath product into $(F_d^{ab} \wr A) \cdot B'$. The first group $F_d^{ab} \wr \mathbb{Z}^n$ is a metabelian group which has finite palindromic width at most $5(d + n)$ by Theorem 3.3. In [AM10] the author proves that every element w of B' is of the form

$$w = [a, t][b, t^2], \quad \text{with } a, b \in B, t \in \mathbb{Z}^n.$$

Since t is an element of an abelian group, it can be written as the product of finitely many palindromes, $t = t_1^{i_1} \cdots t_n^{i_n}$ with $i_j \in \mathbb{Z}$, where $\{t_1, \dots, t_n\}$ is the finite generating set of $\mathbb{Z}^n$. This implies that $t^2 = t_1^{2i_1} \cdots t_n^{2i_n}$. First assume that n is even. Then we can write the first commutator as

$$\begin{aligned} [a, t_1^{i_1} \cdots t_n^{i_n}] &= a^{-1}(t_1^{i_1} \cdots t_n^{i_n})^{-1} a t_1^{i_1} \cdots t_n^{i_n} \\ &= a^{-1} t_n^{-i_n} \overline{a^{-1}} \cdot \overline{a} t_{n-1}^{-i_{n-1}} a \cdot a^{-1} t_{n-2}^{-i_{n-2}} \overline{a^{-1}} \cdots \overline{a} t_1^{-i_1} a \cdot t_1^{i_1} \cdots t_n^{i_n}. \end{aligned}$$

This is now a product of $2n$ palindromes. If n is odd, we write

$$\begin{aligned} [a, t_1^{i_1} \cdots t_n^{i_n}] &= a^{-1}(t_1^{i_1} \cdots t_n^{i_n})^{-1} a t_1^{i_1} \cdots t_n^{i_n} \\ &= a^{-1} t_n^{-i_n} \overline{a^{-1}} \cdot \overline{a} t_{n-1}^{-i_{n-1}} a \cdot a^{-1} t_{n-2}^{-i_{n-2}} \overline{a^{-1}} \cdots a^{-1} t_1^{-i_1} \overline{a^{-1}} \cdot \overline{a} a \cdot t_1^{i_1} \cdots t_n^{i_n}, \end{aligned}$$

which is a product of $2n + 1$ palindromes. We can do the same for the second commutator, hence we need at most $4n$ if n is even and at most $4n + 2$ palindromes if n is odd to express every element of B' . In total we need at most $5(d + n) + 4n = 5d + 9n$ palindromes if n is even and at most $5d + 9n + 2$ palindromes if n is odd. These are palindromes with respect to the natural generating set of $F_d \wr \mathbb{Z}^n$. $\square$

The property of a group of having finite palindromic width with respect to some generating set carries over to its quotients.

Lemma 3.5. *Let G be a group with finite palindromic width $pw(G, X)$ with respect to X . If N is a normal subgroup of G , then $pw(G/N, \hat{X}) \leq pw(G, X)$ where $\hat{X}$ denotes the image of X under the natural homomorphism $G \rightarrow G/N$.*

Every finitely generated group can be obtained as a quotient of some free group. This now implies that Theorem 3.4 in fact holds for any wreath product $G \wr \mathbb{Z}^n$, where G is a finitely generated group.

Corollary 3.6. *The wreath product $G \wr A$ of any d -generated group G with a finitely n -generated abelian group A has finite palindromic width at most $5d + 9n$ if n is even and at most $5d + 9n + 2$ if n is odd with respect to the natural generating set coming from A and G .*

We note however, that this does not imply that $G \wr A$ has finite palindromic width if A is not free abelian. In fact, consider $Q = (G \wr A) / \langle \langle N \rangle \rangle^{G \wr A}$, where $N \triangleleft A$ and $\langle \langle N \rangle \rangle$ denotes the normal closure of N in $G \wr A$. Then for any non-trivial normal subgroup $N \triangleleft A$, the quotient Q becomes metabelian.

If we assume that the group B has finite commutator width and A is an infinite finitely generated abelian group, then it is rather easy to construct palindromes without using the presentation of the commutators from [AM10]. Hence this result also holds if A is not free.

Theorem 3.7. *Let C be a d -generated group with finite commutator width $cw(C) = n$ and A an infinite abelian group of rank r with generating set X such that $|X| = r$. Then*

$$pw(C \wr A, X \cup S) \leq 5d + 6r + 7n$$

for any chosen generating set S of C .

Proof. We write $C \wr A$ as $(C^{ab} \wr A) \cdot (C' \wr A)$. The first factor has finite palindromic width at most $5(d + r)$ by Theorem 3.3. For $C' \wr A$ we do something similar as in the proof of Theorem 4.4. We want to write an element g as a product of finitely many palindromes. Similar to (1), assume g is given as the element

$$g = a \cdot \prod_{i=1}^k a_i^{-1} b_i a_i, \quad b_i = \prod_{j=1}^{n_i} [f_{i,j}, g_{i,j}], \quad n_i \in \mathbb{Z}, b_i, f_i, g_i \in C \times 1.$$

We have to find two elements $s, t \in A$ such that $a_i \neq a_i^{-1}s$ and $a_i \neq a_i^{-1}t \neq a_i^{-1}s$. Since A is infinite abelian this can always be done with s and t both being a different high power of some infinite order generator $x \in X$. Assume $s = x^q$ and $t = x^y$ for some $q, y \in \mathbb{Z}$. We rewrite g with

$$\kappa_j = \prod_{i=1}^k a_{i,j}^{-1} f_{i,j} a_{i,j}, \quad \tau_j = \prod_{i=1}^k a_{i,j}^{-1} g_{i,j} a_{i,j}, \quad h = \prod_{j=1}^n \kappa_j^{-1} \tau_j^{-1} \kappa_j \tau_j$$

as $g = a \cdot h$. We can also write g as

$$g = a \cdot \prod_{j=1}^n \kappa_j^{-1} s^{-1} \overline{\kappa_j^{-1}} \cdot s \cdot \tau_j^{-1} t^{-1} \overline{\tau_j^{-1}} \cdot t \cdot s^{-1} \cdot \overline{\kappa_j} s \kappa_j \cdot t^{-1} \cdot \overline{\tau_j} t \tau_j. \quad (2)$$

The element ts^{-1} is given by x^{y-q} , hence it is a palindrome. We see that this procedure needs at most r palindromes for a and $7n$ palindromes for the product of the commutators in (2). Together we can write every element of $C' \wr H$ under the given hypothesis as a product of at most $r + 7n$ palindromes. With the $5(d + r)$ palindromes from $C^{ab} \wr A$ we hence need at most $5(d + r) + r + 7n = 5d + 6r + 7n$ palindromes to express every element of $C \wr A$. $\square$

4 Wreath Product with a Non-Abelian Group

Let H be a d -generated group and G a group with finite palindromic width with respect to a generating set X . In this Section we show that every element of the subgroup $H' \wr G$ is a product of uniformly bounded finitely many palindromes with respect to a chosen generating set of $H \wr G$.

As a Corollary we deduce that $G \wr K$ has finite palindromic width for all non-abelian finite groups K .

Our main Theorem uses that we can find a word r in the generators of G such that $r = 1$, but $\bar{r} \neq 1$. As we will see in Lemma 4.2, this can be achieved in any non-abelian group of finite palindromic width by eventually adding one element to the generating set. The following example demonstrates that this condition on a word r is easily fulfilled by many groups.

Example 4.1. Consider the group $BS(n, m) = \langle a, b \mid a^{-1}b^na = b^m \rangle$. Clearly $a^{-1}b^na b^{-m} = 1$, but $b^{-m}ab^na^{-1} \neq 1$.

Lemma 4.2. *Assume G is a finitely generated group with finite palindromic width with respect to the generating set X . Then G has a presentation with generating set $X \cup \{c\}$, in which there exists a relation q in G such that $\bar{q} \neq 1$ in G , where $c \in G$ and possibly $c = 1$.*

Proof. Since G has finite palindromic width, it cannot have free quotients. Hence for every pair of non-commuting generators x and y , there exists a relation w , in which x and y occur. Without loss of generality we can assume that x and y occur as subword xy . If there is a subword t between x and y , then either take the first generator used in t that does not commute with x , or otherwise swap x and t in r . So we have

$$r = w_1xyw_2 = 1.$$

If also $\bar{r} = \overline{w_2yxw_1} = 1$, then we introduce a new generator $c = xy$ and get

$$r = w_1cw_2, \quad \bar{r} = \overline{w_2cw_1} = \overline{w_2xyw_1}.$$

Now assume that also under the new generating set $X \cup \{c\}$ we have that $\bar{r} = 1$. This implies

$$\overline{w_2yxw_1} = 1 = \overline{w_2xyw_1},$$

hence x and y commute, a contradiction. $\square$

It is yet unclear how or if the property of having finite palindromic width depends on the chosen generating set. However, as we will see in the following Proposition, enlarging the generating set can only reduce the palindromic width or leave it unchanged.

Proposition 4.3. *Let G be a group which has finite palindromic width with respect to the generating set X . If $Y \subset G$ is any finite subset of G , then $pw(G, X \cup Y) \leq pw(G, X)$.*

Proof. This follows immediately because the set of palindromes in G with respect to $X \cup Y$ contains the set of palindromes of G with respect to X . $\square$

We will now see how we can use this to prove that at every element of the subgroup $F'_d \wr H$ can be expressed as a product of finitely many palindromes.

Theorem 4.4. *Let F_d be a free d -generated group and G a non-abelian group with $pw(G, X) = m$. Then every element of the subgroup $F'_d \wr G$ is a product of at most $m + 1$ palindromes with respect to $X \cup \{c\} \cup Y$, where $c \in G$, possibly $c = 1$.*

The idea of the proof is that the commutator in Figure 1 is a single palindrome of even length and in addition lies in the base group of the wreath product. The elements on the right are the left hand side of the palindrome having positions a_i and the elements on the left are the right hand side of the palindrome on positions $\overline{r^{-1}} \cdot \overline{a_i^{-1}}$ for a relation r that holds in G such that $\bar{r} \neq 1$.




Figure 1: This is how we write one commutator on each position as a palindrome.

Proof of Theorem 4.4. Each element b_i of $F'_d \times 1$ is a product of n_i commutators. Assume that it is the product

$$b_i = \prod_{j=1}^{n_i} [f_{i,j}, g_{i,j}].$$

Now an element g of $F'_d \wr G$ has the form

$$g = a \cdot \prod_{i=1}^k a_i^{-1} b_i a_i = a \cdot \prod_{i=1}^k a_i^{-1} \left(\prod_{j=1}^{n_i} [f_{i,j}, g_{i,j}] \right) a_i, \quad (3)$$

with $f_i, g_i, b_i \in F_d \times 1$. We take into account that we need at most m palindromes for the element a and concentrate on the product of commutators. Take n to be the maximum over all n_i and write every b_i as a product of n (possibly trivial) commutators.

We apply Lemma 4.2 to the generating set of G and choose the relation r in G for which $\bar{r} \neq 1$. We rewrite g with

$$\kappa_j = \prod_{i=1}^n a_i^{-1} f_{i,j}^{-1} r^{-1} g_{i,j}^{-1} r f_{i,j} r^{-1} g_{i,j} r a_i, \quad h = \prod_{i=1}^k \kappa_i,$$

as $g = a \cdot h$. We see that now $\bar{h}$ is given by

$$\bar{h} = \prod_{i=k}^1 \bar{\kappa}_i = \prod_{i=k}^1 \bar{a}_i \cdot \bar{r} \cdot \bar{g}_{i,j} \cdot \bar{r}^{-1} \cdot \bar{f}_{i,j} \cdot \bar{r} \cdot \bar{g}_{i,j}^{-1} \cdot \bar{r}^{-1} \cdot \bar{f}_{i,j}^{-1} \cdot \bar{a}_i^{-1}.$$

We notice that for each i the position of $\bar{f}_{i,j}^{\pm 1}$ is $\bar{a}_i^{-1}$ whereas all $\bar{g}_{i,j}^{\pm 1}$ are at positions $\bar{r}^{-1} \cdot \bar{a}_i^{-1}$. This yields that $\bar{\kappa}_i = 1$ for each $i = 1, \dots, n$ and hence $\bar{h} = 1$. The word $h\bar{h}$ is of course a palindrome and since $\bar{h} = 1$ we have that $h\bar{h} = h$. We form the word

$$w_g = ah\bar{h},$$

which is by construction representing g . We need at most m palindromes to express a and $h\bar{h}$ is exactly one palindrome. Together we can write every element of $F'_d \wr G$ as a product of at most $m+1$ palindromes with respect to the generating set $X \cup \{c\} \cup Y$, with $c \in G$ being possibly trivial. $\square$

We note that if $F_d^{ab} \wr G$ has finite palindromic width for a group G with finite palindromic width, then $F_d \wr G$ has finite palindromic width as well. However, it seems rather difficult to prove that $F_d^{ab} \wr G$ has finite palindromic width in general for any non-abelian group G .

We can again apply Lemma 3.5 and deduce that $H \wr G$ has finite palindromic width for any finitely generated group H and G as in the Theorem above.

Corollary 4.5. *Let H be a finitely generated group and G a non-abelian group with $pw(G, X) = k$. Then every element of the subgroup $H' \wr G$ is a product of at most $k + 1$ palindromes with respect to $X \cup \{c\} \cup Y$, where $c \in G$, possibly trivial.*

A group P is called *perfect* if $P = [P, P]$, in other words, if every element of P is a product of commutators. This can be applied to Theorem 4.4.

Corollary 4.6. *Let P be a finitely generated perfect group and G a non-abelian finitely generated group with finite palindromic width $pw(G, X) = k$ with respect to a generating set X . Then $P \wr G$ has finite palindromic width at most $k + 1$.*

Examples of perfect groups are alternating groups, all simple groups and certain classes of branch groups as described in [LS03] and in a forthcoming paper of this author. It has further been pointed out by D. Gruber that it is possible to construct hyperbolic perfect groups using small cancellation.

If we assume H to be a non-abelian finite group with generating set X , then it follows immediately that every element $h \in H$ is a product of at most $\max \{l_X(h) \mid h \in H\}$ palindromes, where $l_X(h)$ denotes the word length of h with respect to a finite generating set X . With this we can deduce that also the wreath product $F_d \wr H$ has finite palindromic width.

Theorem 4.7. *Assume H is a finite non-abelian group with generating set X and F_d a free d -generated group. Then*

$$pw(F_d \wr H, X \cup \{c\} \cup Y) \leq \max \{l_X(h) \mid h \in H\} \cdot (d \cdot |H| + 1) + 1,$$

with $c \in H$, possibly $c = 1$.

Proof. We split the group $F_d \wr H$ into $(F_d^{ab} \wr H) \cdot (F'_d \wr H)$. The statement for $t \in F'_d \wr H$ follows from Theorem 4.4, using at most $\max \{l_X(h) \mid h \in H\} + 1$ palindromes in the generating set $X \cup \{c\} \cup Y, c \in H$, to express t . An element g of $F_d^{ab} \wr H$ has the form

$$g = a \cdot \prod_{i=1}^{|H|} c_i^{-1} b_i c_i, \quad c_i \in H, b_i \in F_d^{ab} \times 1$$

or more convenient in this case

$$g = \prod_{i=1}^{|H|} a_i b_i, \quad a_i \in H, b_i \in F_d^{ab} \times 1.$$

Every element $b_i \in F_d^{ab} \times 1$ is a product of at most d palindromes and every a_i can be written as a product of at most $\max \{l_X(h) \mid h \in H\}$ palindromes. We do this for every element of H ,

hence we need at most $\max \{l_X(h) \mid h \in H\} \cdot |H| \cdot d$ palindromes for $g \in F'_d \wr H$ and at most $\max \{l_X(h) \mid h \in H\} + 1$ palindromes for $t \in F_d^{ab} \wr H$, which gives at most

$$\max \{l_X(h) \mid h \in H\} \cdot (d \cdot |H| + 1) + 1$$

palindromes for $g \cdot t$. The statement follows since every element of $F_d \wr H$ can be expressed in that way. $\square$

By Lemma 3.5, this also carries over to the wreath product of any finitely generated group G with H as above.

Corollary 4.8. *Assume that H is a non-abelian finite group with generating set X and G a group with d generators. Then*

$$pw(G \wr H, X \cup \{c\} \cup Y) \leq \max \{l_X(h) \mid h \in H\} \cdot (d \cdot |H| + 1) + 1,$$

with $c \in H$, possibly $c = 1$, for any generating set Y of G .

References

- [AM97] M. Akhavan Malayeri. *On the commutator lengths in some finitely generated groups*. Phd-thesis, University of Alberta, 1997.
- [AM06] M. Akhavan-Malayeri. Commutator length of solvable groups satisfying max- n . *Bull. Korean Math. Soc.*, 43(4):805–812, 2006.
- [AM10] M. Akhavan-Malayeri. On commutator length and square length of the wreath product of a group by a finitely generated abelian group. *Algebra Colloq.*, 17(1):799–802, 2010.
- [BG14a] V. Bardakov and K. Gongopadhyay. On palindromic width of certain extensions and quotients of free nilpotent groups. *In Preparation*, 2014.
- [BG14b] V. Bardakov and K. Gongopadhyay. Palindromic width of finitely generated soluble groups. *In Preparation*, 2014.
- [BG14c] V. Bardakov and K. Gongopadhyay. Palindromic width of nilpotent groups. *Journal of Algebra*, 402:379–391, 2014.
- [BST05] V. Bardakov, V. Shpilrain, and V. Tolstykh. On the palindromic and primitive widths of a free group. *Journal of Algebra*, 285(2):574–585, 2005.
- [Col95] D.J. Collins. Palindromic automorphisms of free groups. *London Math. Soc. Lecture Note Ser.*, (204):63–72, 1995.
- [Del10] F. Deloup. Palindromes and orderings in artin groups. *J. Knot Theory Ramifications*, 19(2):145–162, 2010.
- [GC00] H.H. Glover and Jensen C.A. Geometry for palindromic automorphism groups of free groups. *Comment. Math. Helv.*, 75(4):644–667, 2000.

- [LOST10] M.W. Liebeck, E. A. O'Brien, A. Shalev, and P.H. Tiep. The ore conjecture. *J. Eur. Math. Soc. (JEMS)*, 12(4):939–1008, 2010.
- [LOST11] M.W. Liebeck, E.A. O'Brian, A. Shalev, and P.H. Tiep. Commutators in finite quasisimple groups. *Bull. Lond. Math. Soc.*, (6):1079–1092, 2011.
- [LS03] A. Lubotzky and D. Segal. *Subgroup growth*. Number 212 in Progress in Mathematics. Birkhauser Verlag, Basel, 2003.
- [RS14] T. Riley and A. Sale. Palindromic width of wreath products, metabelian groups and solvable max-n groups. <http://arxiv.org/abs/1307.4861>, 2014.
- [Seg09] D. Segal. *Words: notes on verbal width in groups*. Number 361 in London Mathematical Society Lecture Note Series. Cambridge University Press, 2009.